

\$%^STN;HighlightOn= \*\*\*;HighlightOff=\*\*\* ;  
Trying 3106016892...Open

Welcome to STN International! Enter x:x  
LOGINID: ssspta1633cxq  
PASSWORD:  
TERMINAL (ENTER 1, 2, 3, OR 7):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Sep 17 IMSworld Pharmaceutical Company Directory name change  
to PHARMASEARCH  
NEWS 3 Oct 09 Korean abstracts now included in Derwent World Patents  
Index  
NEWS 4 Oct 09 Number of Derwent World Patents Index updates increased  
NEWS 5 Oct 15 Calculated properties now in the REGISTRY/ZREGISTRY File  
NEWS 6 Oct 22 Over 1 million reactions added to CASREACT  
NEWS 7 Oct 22 DGENE GETSIM has been improved  
NEWS 8 Oct 29 AASD no longer available  
NEWS 9 Nov 19 New Search Capabilities USPATFULL and USPAT2  
NEWS 10 Nov 19 TOXCENTER(SM) - new toxicology file now available on STN  
NEWS 11 Nov 29 COPPERLIT now available on STN  
NEWS 12 Nov 29 DWPI revisions to NTIS and US Provisional Numbers  
NEWS 13 Nov 30 Files VETU and VETB to have open access  
NEWS 14 Dec 10 WPINDEX/WPIDS/WPIX New and Revised Manual Codes for  
2002  
NEWS 15 Dec 10 DGENE BLAST Homology Search  
NEWS 16 Dec 17 WELDASEARCH now available on STN  
NEWS 17 Dec 17 STANDARDS now available on STN  
NEWS 18 Dec 17 New fields for DPCL  
NEWS 19 Dec 19 CAS Roles modified  
NEWS 20 Dec 19 1907-1946 data and page images added to CA and Caplus  
NEWS 21 Jan 25 BLAST(R) searching in REGISTRY available in STN on the  
Web  
NEWS 22 Jan 25 Searching with the P indicator for Preparations  
NEWS 23 Jan 29 FSTA has been reloaded and moves to weekly updates  
NEWS 24 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update  
frequency  
NEWS 25 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that  
specific topic.

All use of STN is subject to the provisions of the STN Customer  
agreement. Please note that this agreement limits use to scientific  
research. Use for software development or design or implementation of  
commercial gateways or other similar uses is prohibited and may  
result in loss of user privileges and other penalties.

\*\*\*\*\*STN Columbus\*\*\*\*\*

FILE 'HOME' ENTERED AT 19:15:19 ON 07 MAR 2002

=> FIL BIOSIS EMBASE CAPLUS  
COST IN U.S. DOLLARS SINCE FILE TOTAL  
ENTRY SESSION  
FULL ESTIMATED COST 0.15 0.15

FILE 'BIOSIS' ENTERED AT 19:15:52 ON 07 MAR 2002  
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 19:15:52 ON 07 MAR 2002  
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 19:15:52 ON 07 MAR 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s GPCR (3a) lympho  
L1 0 GPCR (3A) LYMPHO

=> s GPCR (3a) lympho?  
L2 5 GPCR (3A) LYMPHO?

=> dup rem 12  
PROCESSING COMPLETED FOR L2  
L3 3 DUP REM L2 (2 DUPLICATES REMOVED)

=> d bib abs 1-  
YOU HAVE REQUESTED DATA FROM 3 ANSWERS - CONTINUE? Y/(N):y

L3 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2002 ACS  
AN 2000:175932 CAPLUS  
DN 132:218009

TI Protein and cDNA sequences encoding G protein-coupled receptor 14275,  
which is related to the EDG receptor family, and uses thereof in drug  
screening, diagnostic, and therapeutic applications  
IN Glucksmann, Maria Alexandra; Hodge, Martin R.  
PA Millennium Pharmaceuticals, Inc., USA  
SO PCT Int. Appl., 117 pp.  
CODEN: PIXXD2

DT Patent  
LA English  
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PI	WO	2000014233	A1	20000316	WO	1999-US20347	19990903
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM							
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG							
AU	9962434	A1	20000327	AU	1999-62434	19990903	
PRAI	US	1998-146416	A	19980903			
US	1999-390039	A	19990903				
WO	1999-US20347	W	19990903				

AB The invention provides protein and cDNA sequences encoding a novel G protein-coupled receptor (14275), which is a new member of the EDG receptor family. The invention further relates to methods using receptor polypeptides and polynucleotides as a target for diagnosis and treatment in receptor-mediated disorders. The invention further relates to drug-screening methods using the receptor polypeptides and polynucleotides to identify agonists and antagonists for diagnosis and treatment. The invention further encompasses agonists and antagonists based on the receptor polypeptides and polynucleotides.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L3 ANSWER 2 OF 3 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS  
INC.DUPLICATE 1  
AN 2000:48924 BIOSIS  
DN PREV200000048924

TI Kaposi's sarcoma-associated herpesvirus-encoded G protein-coupled receptor activation of c-Jun amino-terminal kinase/stress-activated protein kinase and Lyn kinase is mediated by related adhesion focal tyrosine kinase/proline-rich tyrosine kinase 2.

AU Munshi, Neru; Ganju, Ramesh K.; Avraham, Shalom; Mesri, Enrique A.; Groopman, Jerome E. (1)

CS (1) Division of Experimental Medicine and Hematology/Oncology, Harvard Institutes of Medicine-Beth Israel Deaconess Medical Center, 4 Blackfan Circle, Boston, MA USA

SO Journal of Biological Chemistry, (Nov. 5, 1999) Vol. 274, No. 45, pp. 31863-31867.  
ISSN: 0021-9258.

DT Article  
LA English  
SL English

AB The Kaposi's sarcoma-associated herpesvirus (KSHV) (also known as human herpesvirus 8) has been implicated in the pathogenesis of Kaposi's sarcoma and B cell primary effusion lymphomas. KSHV encodes a G protein-coupled receptor (GPCR) that acts as an oncogene and constitutively activates two protein kinases, c-Jun amino-terminal kinase (JNK)/stress-activated protein kinase (SAPK) and p38 mitogen-activated protein kinase. It also induces the production of vascular endothelial growth factor. These processes are believed to be important in KSHV-GPCR-related oncogenesis. We have characterized the signaling pathways mediated by KSHV-GPCR in a reconstituted 293T cell model in which the related adhesion focal tyrosine kinase (RAFTK) was ectopically expressed. RAFTK has been shown to play an important role in growth factor signaling in endothelium and in B cell antigen receptor signaling in B lymphocytes. KSHV- GPCR induced the tyrosine phosphorylation of RAFTK. Expression of wild-type RAFTK enhanced GPCR-mediated JNK/SAPK activation, whereas dominant-negative mutant constructs of RAFTK, such as K457A (which lacks kinase activity) and Y402F (a Src-binding mutant), inhibited KSHV-GPCR-mediated activation of JNK/SAPK. RAFTK also mediated the KSHV-GPCR-induced activation of Lyn, a Src family kinase. However, RAFTK did not mediate the activation of p38 mitogen-activated protein kinase induced by KSHV-GPCR. Human interferon gamma-inducible protein-10, which is known to inhibit KSHV-GPCR activity, was found to reduce RAFTK phosphorylation and JNK/SAPK activation. These results suggest that in cells expressing RAFTK/proline-rich tyrosine kinase 2, such as endothelial and B cells, RAFTK can act to enhance KSHV-GPCR-mediated downstream signaling to transcriptional regulators such as JNK/SAPK.

L3 ANSWER 3 OF 3 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS  
INC.DUPLICATE 2

AN 1998:406091 BIOSIS  
DN PREV199800406091  
TI Human interferon-gamma-inducible protein 10 (IP-10) inhibits constitutive signaling of Kaposi's sarcoma-associated herpesvirus G protein-coupled receptor.

AU Geras-Raaka, Elizabeth; Varma, Anjali; Ho, Hao; Clark-Lewis, Ian; Gershengorn, Marvin C. (1)  
CS (1) Cornell Univ. Med. Coll., 1300 York Ave., Rm. A328, New York, NY 10021 USA

SO Journal of Experimental Medicine, (July 20, 1998) Vol. 188, No. 2, pp. 405-408.

ISSN: 0022-1007.

DT Article

LA English

AB A G protein-coupled receptor (GPCR) is encoded within the genome of Kaposi's sarcoma-associated herpesvirus (KSHV)/human herpesvirus 8, a virus that may be involved in the pathogenesis of Kaposi's sarcoma and primary effusion \*\*\*lymphomas\*\*\*. KSHV- \*\*\*GPCR\*\*\* exhibits constitutive signaling activity that causes oncogenic transformation. We report that human interferon (IFN)-gamma-inducible protein 10 (HulP-10), a C-X-C chemokine, specifically inhibits signaling of KSHV-GPCR. In contrast, monokine induced by IFN-gamma (HuMig), which like HulP-10 is an agonist of C-X-C chemokine receptor 3, does not inhibit KSHV-GPCR signaling. Moreover, HulP-10, but not HuMig, inhibits KSHV-GPCR-induced proliferation of NIH 3T3 cells. These results show that HulP-10 is an inverse agonist that converts KSHV-GPCR from an active to an inactive state. Thus, a human chemokine inhibits constitutive signaling and cellular proliferation that is mediated by a receptor encoded by a human disease-associated herpesvirus.

=> s GPCR (5a) lympho?

L4 6 GPCR (5A) LYMPHO?

=> dup rem l4

PROCESSING COMPLETED FOR L4

L5 4 DUP REM L4 (2 DUPLICATES REMOVED)

=> d bib abs 1-

YOU HAVE REQUESTED DATA FROM 4 ANSWERS - CONTINUE? Y/(N):y

L5 ANSWER 1 OF 4 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

AN 2001:289395 BIOSIS

DN PREV200100289395

TI Refolding and isolation of an intact SLC6Kine expressed as an insoluble fusion protein in E coli.

AU Johanson, Kyung O. (1); Scott, Gilbert (1); McDevitt, Patrick (1); Matico, Rosalie (1); Sarau, Henry (1); Appelbaum, Edward (1)

CS (1) Protein Biochemistry, SmithKline Beecham Pharm, 709 Swedeland Rd, King of Prussia, PA, 19406 USA

SO FASEB Journal, (March 7, 2001) Vol. 15, No. 4, pp. A530. print.  
Meeting Info: Annual Meeting of the Federation of American Societies for Experimental Biology on Experimental Biology 2001 Orlando, Florida, USA March 31-April 04, 2001  
ISSN: 0892-6638.

DT Conference

LA English

SL English

AB SLC (secondary lymphoid tissue chemokine) is a C-C chemokine with three disulfide bonds (6CKine) in 111 amino acids. It is a potent chemotactic factor for dendritic cells and T \*\*\*lymphocytes\*\*\* mediated through a 7TM \*\*\*GPCR\*\*\* receptor, CCR7. To investigate its potential as a drug target, various expression systems were tried in order to produce sufficient amount of an intact molecule. Natural N-terminus was determined using Drosophila secretion system using its own signal sequence, however the expression level was very low. The protein was then expressed at high level in E. coli as an insoluble protein with an N-terminal 6His tag with a cleavage site of enterokinase (EK) or factor Xa, or Genenase. The inclusion body was solubilized and purified using NiNTA agarose in 6 M GdnHCl and reduced with DTT. GdnHCl and DTT were removed from the reduced/denatured chemokine by dialyzing against 5 mM HCl. It was then refolded by air oxidation at pH 8.5. EK or Xa treatment resulted in multiple cleavages at C-terminal region, while Genenase yielded an intact molecule. The chemokine was active in calcium mobilization assays with cells expressing recombinant CCR7 (EC50 = 1.2 nM).

L5 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2002 ACS

AN 2000:175932 CAPLUS

DN 132:218009

TI Protein and cDNA sequences encoding G protein-coupled receptor 14275, which is related to the EDG receptor family, and uses thereof in drug screening, diagnostic, and therapeutic applications

IN Glucksmann, Maria Alexandra; Hodge, Martin R.

PA Millennium Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 117 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN,CNT 1

PATENT NO. KIND DATE APPLICATION NO. DATE

PI WO 2000014233 A1 20000316 WO 1999-US20347 19990903

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 9962434 A1 20000327 AU 1999-62434 19990903

PRAI US 1998-146416 A 19980903

US 1999-390039 A 19990903

WO 1999-US20347 W 19990903

AB The invention provides protein and cDNA sequences encoding a novel G protein-coupled receptor (14275), which is a new member of the EDG receptor family. The invention further relates to methods using receptor polypeptides and polynucleotides as a target for diagnosis and treatment in receptor-mediated disorders. The invention further relates to drug-screening methods using the receptor polypeptides and polynucleotides to identify agonists and antagonists for diagnosis and treatment. The invention further encompasses agonists and antagonists based on the receptor polypeptides and polynucleotides.

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 3 OF 4 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE 1

AN 2000:48924 BIOSIS

DN PREV20000048924

TI Kaposi's sarcoma-associated herpesvirus-encoded G protein-coupled receptor activation of c-Jun amino-terminal kinase/stress-activated protein kinase and Lyn kinase is mediated by related adhesion focal tyrosine kinase/proline-rich tyrosine kinase 2.

AU Munshi, Neru; Ganju, Ramesh K.; Avraham, Shalom; Mesri, Enrique A.; Groopman, Jerome E. (1)

CS (1) Division of Experimental Medicine and Hematology/Oncology, Harvard Institutes of Medicine-Beth Israel Deaconess Medical Center, 4 Blackfan Circle, Boston, MA USA

SO Journal of Biological Chemistry, (Nov. 5, 1999) Vol. 274, No. 45, pp. 31863-31867.

ISSN: 0021-9258.

DT Article

LA English

SL English

AB The Kaposi's sarcoma-associated herpesvirus (KSHV) (also known as human herpesvirus 8) has been implicated in the pathogenesis of Kaposi's sarcoma and B cell primary effusion lymphomas. KSHV encodes a G protein-coupled receptor (GPCR) that acts as an oncogene and constitutively activates two protein kinases, c-Jun amino-terminal kinase (JNK)/stress-activated protein kinase (SAPK) and p38 mitogen-activated protein kinase. It also induces the production of vascular endothelial growth factor. These processes are believed to be important in KSHV-GPCR-related oncogenesis. We have characterized the signaling pathways mediated by KSHV-GPCR in a reconstituted 293T cell model in which the related adhesion focal tyrosine kinase (RAFTK) was ectopically expressed. RAFTK has been shown to play an important role in growth factor signaling in endothelium and in B cell antigen receptor signaling in B \*\*\*lymphocytes\*\*\*. KSHV- \*\*\*GPCR\*\*\* induced the tyrosine phosphorylation of RAFTK. Expression of wild-type RAFTK enhanced GPCR-mediated JNK/SAPK activation, whereas dominant-negative mutant constructs of RAFTK, such as K457A (which lacks kinase activity) and Y402F (a Src-binding mutant), inhibited KSHV-GPCR-mediated activation of JNK/SAPK. RAFTK also mediated the KSHV-GPCR-induced activation of Lyn, a Src family kinase. However, RAFTK did not mediate the activation of p38 mitogen-activated protein kinase induced by KSHV-GPCR. Human interferon gamma-inducible protein-10, which is known to inhibit KSHV-GPCR activity, was found to reduce RAFTK phosphorylation and JNK/SAPK activation. These results suggest that in cells expressing RAFTK/proline-rich tyrosine kinase 2, such as endothelial and B cells, RAFTK can act to enhance KSHV-GPCR-mediated downstream signaling to transcriptional regulators such as JNK/SAPK.

L5 ANSWER 4 OF 4 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE 2

AN 1998:406091 BIOSIS

DN PREV199800406091

TI Human interferon-gamma-inducible protein 10 (IP-10) inhibits constitutive signaling of Kaposi's sarcoma-associated herpesvirus G protein-coupled receptor.

AU Geras-Raaka, Elizabeth; Varma, Anjali; Ho, Hao; Clark-Lewis, Ian; Gershengorn, Marvin C. (1)

CS (1) Cornell Univ. Med. Coll., 1300 York Ave., Rm. A328, New York, NY 10021 USA

SO Journal of Experimental Medicine, (July 20, 1998) Vol. 188, No. 2, pp. 405-408.

ISSN: 0022-1007.

DT Article

LA English

AB A G protein-coupled receptor (GPCR) is encoded within the genome of Kaposi's sarcoma-associated herpesvirus (KSHV)/human herpesvirus 8, a virus that may be involved in the pathogenesis of Kaposi's sarcoma and primary effusion \*\*\*lymphomas\*\*\*. KSHV- \*\*\*GPCR\*\*\* exhibits

constitutive signaling activity that causes oncogenic transformation. We report that human interferon (IFN)-gamma-inducible protein 10 (HulP-10), a C-X-C chemokine, specifically inhibits signaling of KSHV-GPCR. In contrast, monokine induced by IFN-gamma (HuMig), which like HulP-10 is an agonist of C-X-C chemokine receptor 3, does not inhibit KSHV-GPCR signaling. Moreover, HulP-10, but not HuMig, inhibits KSHV-GPCR-induced proliferation of NIH 3T3 cells. These results show that HulP-10 is an inverse agonist that converts KSHV-GPCR from an active to an inactive state. Thus, a human chemokine inhibits constitutive signaling and cellular proliferation that is mediated by a receptor encoded by a human disease-associated herpesvirus.

=>

---Logging off of STN---

=>

Executing the logoff script..

=> LOG Y